

## Myofascial release; an evidence based treatment concept?

Lars Remvig, MD, DMSc

Department of Medical Orthopaedics and Rehabilitation, Rigshospitalet, University Hospital of Copenhagen, Blegdamsvej 9, 2100 Copenhagen, Denmark.

phone +45 3545 2005, fax, +45 3545 2008, email [lars.remvig@rh.regionh.dk](mailto:lars.remvig@rh.regionh.dk)

**BACKGROUND** Myofascial release (MFR) is a very commonly used treatment modality for painful somatic dysfunctions. However, do we have any scientific evidence for an effect of MFR?

**METHODS** A literature search is performed in order to elucidate the prerequisites for myofascial pain, the interobserver variability of diagnostic tests for myofascial dysfunction and the clinical effect of MFR.

**RESULTS** We have evidence that fascia are built by stretchable collagenous elements, that fascia contains contractile elements, that fascia have vascular and neural supply, and that the dermis is attached to the underlying fascia.

We have no studies that describe intertester reproducibility of diagnostic tests for fascial tightness or looseness and therefore we cannot say for sure that either situation can generate pain. Neither is there any studies documenting that tight fascia can be released – i.e stretched – by MFR technique or that the technique *per se* gives relief to the patients myofascial pain.

**CONCLUSION** The prerequisites for myofascial pain is present. However, the proof for the MFR clinical diagnosis and the effect of the MFR treatment is lacking.

### REFERENCE

[1] Staubesand J, Li Y. Zum fainbau der fascia cruris mit besonderer berücksichtigung epi- und intrafascialen nerven. Manuelle Medizin 1996;34:196-200.

[2] Hinz B, Celetta G, Tomasek JJ, Gabbiani G, Chaponnier C. Alpha-smooth muscle actin expression upregulates fibroblast contractile activity. Mol Biol Cell 2001;12:2730-41.

[3] Schleip R. Active fascial contractility. Implications for musculoskeletal mechanics. Thesis. Faculty of Medicine, Ulm University, Germany. 2006