

# How Could We Explain the Changes of the Plantar Muscle Tone in the Patients with Type I Diabetes Mellitus and Interpret These Findings

MD, PhD Mr. R. Kerpe, MD, PhD Mr. A. Krisciunas

Department of Rehabilitation , Kaunas University of Medicine, Eiveniu Str. 2, LT- 44307, Kaunas, Lithuania e-mail: [riciricardo@delfi.lt](mailto:riciricardo@delfi.lt)

**PURPOSE:** The investigators determined that thickening of the *plantar fascia* in diabetics concurs to develop a rigid foot, which poorly absorbs shock during loading. They concluded that the diabetic foot often undergoes abnormal plantar pressures, changing walking strategy and ulcerative processes [1; 2]. In our study, we investigated the muscle tone in patients with type I diabetes mellitus (DM). We hypothesized that the muscle tone in the foot could be partly influenced by the changes in the *extracellular matrix*.

**METHODS:** The study involved randomly selected 70 subjects with type 1 diabetes mellitus: 31 men (44.3%) and 39 women (55.7%) at the age of  $32.4 \pm 10.3$  years. The duration of diabetes was  $18.7 \pm 8.3$  years; it also involved randomly selected 31 healthy subjects as controls – 12 men (38.7%) and 19 women (61.3%) at the age of  $33.3 \pm 10.0$  years. Assessment of the selected patients was performed using a standardized questionnaire and examination. An original method (myotonometry) and equipment, which makes it possible to measure the parameters of skeletal muscles, was used in this study.

**RESULTS:** It was estimated that the mean of the tone of the *flexor digitorum brevis* muscle in the patients with type 1 diabetes mellitus was increased compared with healthy controls.

**CONCLUSIONS:** In 2008, *Bus Sicco A* presented a publication in which he postulated that patients with DM could have more thickened *plantar aponeurosis*. This thickening contributes to a more rigid *cavus* like foot that bears increased forefoot loads during foot progression [3]. This supposition correlates with *Elhadd TA et al.* publication in which the thickening of the *plantar aponeurosis* in association with Dupuytren's disease in the patient with DM was considered [4]. These two publications, together with the *Schleip R.* publication (Schleip R., 2006) on the possibility to influence the myofibroblast activity and physical properties of the connective tissue, enabled us to make the following conclusions:

1. Diabetes mellitus can cause thickening of the *plantar aponeurosis* similar to the one in Dupuytren's contracture. The thickening is a result of the increased activity of fibroblasts.
2. The therapeutic application of the rehabilitation methods influencing fibroblast activity could prevent (postpone) the development of the thickened *plantar aponeurosis* and provide a normal development (especially in children and adolescents) of foot architecture in diabetes mellitus.

## REFERENCES:

1. D'Ambrogi ED, Giurato L, D'Agostino MA, Giacomozzi C, Macellari V, Caselli A, Uccioli L. Contribution of plantar fascia to the increased forefoot pressures in diabetic patients. *Diabetes Care* 2003; 26:1525–1529.
2. Giacomozzi C, D'Ambrogi E, Uccioli L, Macellari V. Does the thickening of Achilles tendon and plantar fascia contribute to the alteration of diabetic foot loading? *Clin Biomech* (Bristol, Avon) 2005 Jun;20(5):532-9.
3. Bus Sicco A. Foot structure and footwear prescription in diabetes mellitus. *Diabetes Metab Res Rev* 2008; 24(Supp 1):S90-S95.
4. Elhadd TA, Ghosh S, Malik MI, Collier A. Plantar fibromatosis and Dupuytren's disease: an association to remember in patients with diabetes. (Letter) *Diabet Med* 2007 Nov;24(11): 1305.
5. Schleip R, Naylor IL, Ursu D, Melzer W, Zorn A, Wilke HJ, Lehmann-Horn F, Klingler W. Passive muscle stiffness may be influenced by active contractility of intramuscular connective tissue. *Med Hypotheses* 2006; 66(1):66–71.