

# The Strain Patterns of the Deep Fascia of the Lower Limb


Andrew Franklyn-Miller<sup>1</sup>, Eanna Falvey<sup>1</sup>, Ross Clark<sup>1</sup>, Adam Bryant<sup>1</sup>, Peter Brukner<sup>1</sup>, Priscilla Barker<sup>2</sup>, Christopher Briggs<sup>2</sup> Paul McCrory<sup>1</sup> Email [afranklynmiller@me.com](mailto:afranklynmiller@me.com)<sup>1</sup> Centre for Health, Exercise and Sports Medicine, Faculty of Medicine, Dentistry and Health Sciences, 2 Department of Anatomy and Cell Biology, Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne, Victoria 3010 Australia.

**BACKGROUND** There are many theories as to lower limb fascial function including the hydraulic amplifier effect<sup>1</sup> with respect to lumbar fascia, the crossover effect in support of bipedal motion<sup>2</sup> and tensioning theory<sup>3</sup>. Schleip, et al<sup>4</sup> have demonstrated contractile properties of fascia and Barker<sup>5</sup> and others have demonstrated discrete collagen fiber orientation. The authors set out to find patterns of strain in the lower limb fascia when performing a movement protocol

**METHODS** Five, fresh cadavers were used. Insulated, 10mm, 120Ohm foil-type micro-strain gauges (BCM Sensor Technologies, Belgium) were attached to the external surface of the fascia (Fig.1) using a gauge specific cyanoacrylate adhesive (TML, Japan) prior to the performance of a movement protocol of the lower limb. Data was acquired at 50Hz via a USB-based CompactDAQ system, and was normalized and calibrated using a combination of Signal Express 2.0 and Labview 8.5 software (National Instruments, U.S.A.).

**RESULTS** Microstrain(me) results for the straight leg raise (SLR) are presented below, strain gauges were parallel to gross fibre direction.

Location of Strain Gauge	Median Strain in SLR
Contralateral Lumbar Fascia	6.7
Ipsilateral Lumbar Fascia	21.8
ITB	35.3
Posterior Thigh	14.9
Lateral Compartment	15.3
Achilles	15.0
Plantar Fascia	3.9



**CONCLUSIONS** This work concurs with the theory of fascial strain transmission in the lumbar and lower limb fascia and with collagen fibre orientation as per earlier work provides potential insight into the fascial role in overuse musculoskeletal disorders. Fig 1. Micro-strain gauges on lumbar fascia

1. Aspden RM. Review of the functional anatomy of the spinal ligaments and the lumbar erector spinae muscles. *Clinical Anatomy* 1992(5):372-87.
2. Vleeming A, Pool-Goudzwaard AL, Stoeckart R, van Wingerden JP, Snijders CJ. The posterior layer of the thoracolumbar fascia. Its function in load transfer from spine to legs. *Spine* 1995;20(7):753-8.
3. Barker PJ, Briggs CA, Bogeski G. Tensile transmission across the lumbar fasciae in unembalmed cadavers: effects of tension to various muscular attachments. *Spine* 2004;29(2):129-38.
4. Active contraction of the thoracolumbar fascia - indications of a new factor in low back pain research with implications for manual therapy. Fifth Interdisciplinary World Congress of Low Back Pain and Pelvic Pain; 2004; Melbourne, Australia.
5. Barker PJ, Briggs CA. Attachments of the posterior layer of lumbar fascia. *Spine* 1999;24(17):1757-64.